

# **Tobacco Prevention and Control in WA**

University of Washington April 18, 2013

Paul Davis, Manager, Tobacco Prevention and Control Unit
Washington State Department of Health



### **Asthma Rates for Kids and Adults**

Figure 14. Youth with lifetime asthma: Washington, 2002-2010







Figure 2. Adults with current asthma: Washington and U.S., 1999-2010





#### **Cancer Incidence and Death**







PUBLIC HEALTH
ALWAYS WORKING FOR A SAFER AND
HEALTHIER WASHINGTON



### **Heart Disease and Diabetes**








#### Four Goals of Tobacco Prevention and Control

- 1. Identify and reduce tobacco related disparities
- 2. Prevent youth from starting to use tobacco
- 3. Help people quit using tobacco
- 4. Eliminate exposure to secondhand smoke





## **Best Practice Programs Take Money but...**








# Federal, State, and Local Funding for Tobacco

- CDC provides \$1.7 million/yr in core tobacco funds
- Community Transformation Grant (CTG)
  - WA state grant
  - King, Pierce, Spokane grants
  - Chehalis and Makah tribes
- Youth Tobacco Prevention \$750k/yr from licenses and fines
- Some counties receive local funding






## **Tobacco Taxes Have Increased Over Time**



# Sales of Tobacco to Minors




**Washington State** 



Source: Washington State Department of Health Synar Compliance Check Database



# **Cigarette Smoking Among Youth**




After what seemed like a flattening of the trend from 2004-2010, all grades showed decreases from 2010 to 2012

ALWAYS WORKING FOR A SAFER AND HEALTHIER WASHINGTON

10



## 10th Grade Youth Tobacco Rates











# **Pregnancy and Smoking**



- Smoking 3 months before pregnancy
- Smoking last 3 months of pregnancy
- → Post Partum Smoking








\* Data are from the Behavioral Risk Factor Surveillance System.

NOTE: Lower Income was defined as a yearly household income of less \$25,000.

Lower Education was defined as a high school degree or less among respondents 25 and older.

## Smoking Among Adults by Sex and Age in Washington, 2011



Source: Washington Behavioral Risk Factor Survey \* Age adjusted percent.

\*\* Insufficient data

